arXiv:1704.06715v1 [math.CO] 21 Apr 2017

# Face enumeration on matroid base polytopes

Vladimir Grujić
Faculty of Mathematics
Belgrade University
Serbia
vgrujic@matf.bg.ac.rs

Marko Pešović
Faculty of Civil Engineering
Belgrade University
Serbia
mpesovic@grf.bg.ac.rs

Mathematics Subject Classifications: 52B40, 52B05, 16T05

**Abstract**

We introduce a refinement of the Billera, Jia and Reiner quasisymmetric function that carries information of face numbers of generalized permutohedra. This includes a refinement of Stanley's chromatic symmetric function. We consider more systematically the case of matroid base polytopes and obtain the formula for their $f$-polynomials in terms of principal specializations of quasisymmetric enumerators.

**Keywords**: generalized permutohedron, quasisymmetric function, matroid, matroid base polytope, combinatorial Hopf algebra, f-polynomial

## 1 Introduction

For a generalized permutohedron $Q$ there is a quasisymmetric enumerator function $F(Q)$ introduced by Billera, Jia and Reiner in [3]. It enumerates positive integer lattice points $\omega = (\omega_1, \ldots, \omega_n) \in \mathbb{Z}_+^n$ which are $Q$-generic. It means that the weight function $\omega^\star : Q \to \mathbb{R}$ defined by $\omega^\star(v) = \langle \omega, x \rangle$ has its maximum at a unique vertex $v$ of $Q$. That is

$$F(Q) = \sum_{\omega \ Q-\text{generic}} \mathbf{x}_\omega, \tag{1}$$

where $\mathbf{x}_\omega = x_{\omega_1} x_{\omega_2} \cdots x_{\omega_n}$. The generalized permutohedra are introduced and extensively studied by Postnikov [15] and Postnikov, Reiner and Williams [16]. They are deformations of the standard permutohedra $Pe^{n-1}$. A part of the reach combinatorial structure of a generalized permutohedron $Q$ is a certain statistic $\mathrm{rk}_Q(\mathcal{F})$ on the face lattice $L(Pe^{n-1})$ of the standard permutohedron $Pe^{n-1}$ which we call the *Q-rank*. The space $\mathbb{R}^n$ is decomposed by faces of the braid arrangement fan $\Sigma_{Pe^{n-1}}$ which is the normal fan of $Pe^{n-1}$. Each vector $\omega \in \mathbb{Z}_+^n$ lies in the relative interior of the normal cone at a unique face $\mathcal{F}_\omega$ of $Pe^{n-1}$. We refine the Billera, Jia and Reiner quasisymmetric invariant by

$$F_q(Q) = \sum_{\omega \in \mathbb{Z}_+^n} q^{\mathrm{rk}_Q(\mathcal{F}_\omega)} \mathbf{x}_\omega. \tag{2}$$

The $Q$-rank of a face $\mathcal{F}$ is zero only for vertices, so $F_0(Q) = F(Q)$.

On the other hand, when a class of generalized permutohedra is specified we obtain the known combinatorial enumerators. The case of graphical zonotopes $Q = Z_\Gamma$ is studied in [11]. The quasisymmetric function $F_q(Z_\Gamma)$ is a $q$-refinement of the Stanley chromatic symmetric function of a graph $X_\Gamma$. It is obtained as a specialization at $q = 0$

$$F_0(Z_\Gamma) = X_\Gamma.$$

The interesting results are also obtained in the case of nestohedra studied in [10] and for a $q$-analog in [12]. For a subclass of graph-associahedra $Q = P_\Gamma$ the enumerator $F(P_\Gamma)$ produces a new quasisymmetric invariant of a graph $\Gamma$.

The both cases of graphical zonotopes and nestohedra have in common that the enumerator $F(Q)$ and its $q$-analog $F_q(Q)$ coincide with the universal morphism from certain combinatorial Hopf algebras to quasisymmetric functions. In the case of graphical zonotopes it is the chromatic Hopf algebra of graphs and in the case of nestohedra it is the non-cocommutative Hopf algebra on building sets and its Hopf subalgebra of graphs.

Billera, Jia and Reiner applied their enumerator $F(Q)$ in the case of matroid base polytopes $Q = P_M$. The invariant $F(P_M)$ also comes from a Hopf algebra, in this case of matroids which was firstly introduced by Schmitt [17].

The paper is organized as follows. In section 2 we review the necessary facts about combinatorics of standard permutohedron and introduce

generalized permutohedra. In section 3 we review the basic facts about quasisymmetric functions. In section 4 we introduce the quasisymmetric function $F_q(Q)$ and show that it contains the information about $f$-vectors of generalized permutohedra which provides a unifying perspective on the main results of the papers [11] and [12]. It also may be regarded as a far-reaching illumination of the Stanley $(-1)$-color theorem for numbers of acyclic orientations of a graph. In the rest of the paper the case of matroid base polytopes is considered more thoroughly. In section 5 we review some basic facts about combinatorics of matroid base polytopes. The combinatorial Hopf algebra on matroids and its $q$-analog are introduced. Finally some calculations concerning the case of uniform matroids is presented.

# 2 Generalized permutohedra

The symmetric group $S_n$ acts on the space $\mathbb{R}^n$ by permuting the coordinates. Recall that a $(n-1)$-dimensional *permutohedron* $Pe^{n-1}$ is the convex hull of the orbit of a point with increasing coordinates $x_1 < \cdots < x_n$

$$Pe^{n-1} = \mathrm{Conv}\{(x_{\omega(1)}, x_{\omega(2)} \ldots, x_{\omega(n)}) \mid \omega \in S_n\}.$$

The *braid arrangement* is the arrangement of hyperplanes $\{x_i = x_j\}_{1 \le i < j \le n}$ in the space $\mathbb{R}^n$. The regions of braid arrangement, called *Weyl chambers* are labelled by permutations $\omega \in S_n$

$$C_\omega := \{x_{\omega(1)} \le x_{\omega(2)} \le \ldots \le x_{\omega(n)}\}.$$

The corresponding *braid arrangement fan* is the normal fan $\Sigma_{Pe^{n-1}}$ of the permutohedron $Pe^{n-1}$.

A flag of the length $|\mathcal{F}| = k$ on the set $[n] = \{1, \ldots, n\}$ is a chain of subsets

$$\mathcal{F} : \emptyset =: F_0 \subset F_1 \subset \ldots \subset F_{k-1} \subset F_k \subset F_{k+1} := [n].$$

The type of a flag $\mathcal{F}$ is the following $(k+1)$-tuple of integers

$$\mathrm{type}(\mathcal{F}) = (|F_1| - |F_0|, |F_2| - |F_1|, \ldots, |F_{k+1}| - |F_k|).$$

The set of flags is ordered by refinements. There is an obvious order reversing one-to-one correspondence between the face lattice of the permutohedron $Pe^{n-1}$ and the lattice of flags of $[n]$. With no abuse of notation we denote

a face of the permutohedron $Pe^{n-1}$ by the corresponding flag $\mathcal{F}$. By this convention we have that $\dim\mathcal{F} = n - |\mathcal{F}| - 1$. For example, the facets correspond to the flags $\emptyset \subset A \subset [n]$, while the vertices correspond to maximal flags.

The normal cone $\sigma_{\mathcal{F}} \in \Sigma_{Pe^{n-1}}$ at the face $\mathcal{F}$ is determined by the coordinates relations

1. $x_p = x_q$ if $p, q \in F_{i+1} \setminus F_i$ for some $i = 0, \ldots, k$,
2. $x_p \leq x_q$ if $p \in F_i \setminus F_{i-1}$ and $q \in F_{i+1} \setminus F_i$ for some $i = 1, \ldots, k$.

Note that $\dim(\sigma_{\mathcal{F}}) = |\mathcal{F}|$ and the relative interior $\sigma^\circ_{\mathcal{F}}$, given by strict inequalities in the condition 2 above, is homeomorphic to $\mathbb{R}^{|\mathcal{F}|}$. Conversely, the flag $\mathcal{F}$ can be reconstructed from the cone $\sigma_{\mathcal{F}}$ by setting $p \in F_{i+1} \setminus F_i$ and $q \in F_{j+1} \setminus F_j$ for some $0 \leq i < j \leq k$ whenever $x_p < x_q$ for all points in $\sigma^\circ_{\mathcal{F}}$.

**Definition 2.1.** A convex polytope $Q$ is an $(n-1)$-dimensional *generalized permutohedron* if its normal fan $\Sigma_Q$ is coarser than the braid arrangement fan $\Sigma_{Pe^{n-1}}$.

Generalized permutohedra are equivalently characterized as deformations of the standard permutohedron $Pe^{n-1}$ by moving its vertices with keeping directions of edges. Thus any edge of a generalized permutohedron $Q$ lies in the direction of some $e_i - e_j$, where $e_i, i = 1, \ldots, n$ are the standard basis vectors in $\mathbb{R}^n$.

For an $(n-1)$-dimensional generalized permutohedron $Q$ is defined the map

$$\pi_Q : L(Pe^{n-1}) \to L(Q)$$

of face lattices by $\pi_Q(\mathcal{F}) = G$ if and only if the relative interior $\sigma^\circ_{\mathcal{F}}$ of the normal cone at the face $\mathcal{F}$ of $Pe^{n-1}$ is contained in the relative interior $\sigma^\circ_G$ of the normal cone at the face $G$ of $Q$.

**Proposition 2.2.**

$$\sum_{\mathcal{F}:\pi_Q(\mathcal{F})=G} (-1)^{|\mathcal{F}|} = (-1)^{n-\dim(G)-1}.$$

*Proof.* Consider the collection of flags $\pi_Q^{-1}(G) = \{\mathcal{F}_1, \mathcal{F}_2, \ldots \mathcal{F}_t\}$. We have $\sigma^\circ_G = \bigcup_{i=1}^t \sigma^\circ_{\mathcal{F}_i}$ and $\sigma^\circ_G$ is homeomorphic to the $k$-dimensional open cell, where

$k = n - \dim G - 1$. Let $f_i, i = 0, 1, \ldots, k$ be the numbers of $i$-dimensional cones such that $\sigma^\circ_{\mathcal{F}} \subset \sigma^\circ_G$. By inclusion-exclusion principle we have

$$f_k - f_{k-1} + \cdots + (-1)^k f_0 = 1.$$

□

# 3 Quasisymmetric functions

In this section we review the basic facts about combinatorial Hopf algebras and quasisymmetric functions. The notion of combinatorial Hopf algebra is well founded in the work of Aguiar, Bergeron and Sottile [1]. The extensive survey of Hopf algebra theory in combinatorics may be found in [9].

A combinatorial Hopf algebra is a graded, connected Hopf algebra $\mathcal{H}$ equipped with a multiplicative linear functional to the ground field $\zeta : \mathcal{H} \to \mathbf{k}$. We describe the combinatorial Hopf algebra $QSym$ of quasisymmetric functions which is the terminal object in the category of combinatorial Hopf algebras.

A quasisymmetric function $F = F(\mathbf{x})$ is a formal power series of bounded degree in the countable ordered set of variables $\mathbf{x} = (x_1, x_2, \ldots)$ such that coefficients by monomials with the same list of ordered exponents are equal. This condition produces the natural linear basis for the algebra $QSym$ consisting of monomial quasisymmetric functions

$$M_\alpha = \sum_{i_1 < i_2 < \cdots < i_k} x_{i_1}^{a_1} x_{i_2}^{a_2} \cdots x_{i_k}^{a_k},$$

indexed by finite ordered sets of integers $\alpha = (a_1, a_2, \ldots, a_k)$ called compositions of $|\alpha| = a_1 + a_2 + \cdots a_k$. The coproduct, defined on monomial basis by

$$\Delta(M_\alpha) = \sum_{\beta\gamma = \alpha} M_\beta \otimes M_\gamma,$$

where $\beta\gamma$ is the concatenation of compositions, turns $QSym$ into a graded, connected Hopf algebra.

Another standard linear basis of $QSym$ is consisting of fundamental quasisymmetric functions $L_\alpha$. A composition $\alpha = (a_1, a_2, \ldots, a_k)$ of the length $k(\alpha) = k$ is uniquely determined by the $(k-1)$-elements subset

$D(\alpha) = \{a_1, a_1 + a_2, \ldots, a_1 + \cdots a_{k-1}\}$ of the set $[|\alpha|]$. The set of compositions of an integer $n$ is ordered by the refinement relation defined by $\alpha \preceq \beta$ if and only if $D(\alpha) \subseteq D(\beta)$. Then

$$L_\alpha = \sum_{\alpha \preceq \beta} M_\beta.$$

The principal specialization $\mathbf{ps} : QSym \to \mathbf{k}[m]$ assigns to a quasisymmetric function $F$ a polynomial in $m$ by evaluation map

$$\mathbf{ps}(F)(m) = F|_{x_1=\cdots=x_m=1, x_{m+1}=\cdots=0}.$$

The canonical character on quasisymmetric functions $\zeta_Q : QSym \to \mathbf{k}$ is defined by $\zeta_Q(F) = \mathbf{ps}(F)(1)$. It is easy to see that on monomial basis we have

$$\mathbf{ps}(M_\alpha)(m) = \binom{m}{k(\alpha)}$$

and specially

$$\mathbf{ps}(M_\alpha)(-1) = (-1)^{k(\alpha)}.$$

The importance of the antipode $S : \mathcal{H} \to \mathcal{H}$ in enumerative problems is indicated by

$$\mathbf{ps}(F)(-1) = \zeta \circ S(h).$$

The following theorem is fundamental in applications and explains the ubiquity of quasisymmetric functions in combinatorics.

**Theorem 3.1** ([1], Theorem 4.1)**.** *For a combinatorial Hopf algebra $(\mathcal{H}, \zeta)$ there is a unique morphism of graded Hopf algebras*

$$\Psi : \mathcal{H} \to QSym$$

*such that $\Psi \circ \zeta_Q = \zeta$. The coefficients $c_\alpha(h)$ in monomial basis of $\Psi(h)$ for a homogeneous element $h$ of degree $n$ are given by*

$$c_\alpha(h) = \zeta^{\otimes k} \circ (p_{a_1} \otimes p_{a_2} \otimes \cdots \otimes p_{a_k}) \circ \Delta^{(k-1)}(h),$$

*where $p_i$ is the projection on the $i$-th homogeneous component and $\Delta^{(k-1)}$ is the $(k-1)$-fold coproduct map of $\mathcal{H}$.*

# 4 Quasisymmetric enumerator $F_q(Q)$

Recall that to a generalized permutohedron $Q$ is assigned the map $\pi_Q : L(Pe^{n-1}) \to L(Q)$ between face lattices by $\pi_Q(\mathcal{F}) = G$ if and only if $\sigma^\circ_{\mathcal{F}} \subset \sigma^\circ_G$. We derive a natural statistic of faces of the standard permutohedron $Pe^{n-1}$ determined by $Q$.

**Definition 4.1.** For a generalized permutohedron $Q$ the $Q$-*rank* is a map on the face lattice of the standard permutohedron $\mathrm{rk}_Q : L(Pe^{n-1}) \to \{0, 1, \ldots, n-1\}$ defined by

$$\mathrm{rk}_Q(\mathcal{F}) = \dim(\pi_Q(\mathcal{F})).$$

Let $\omega = (\omega_1, \omega_2, \ldots, \omega_n) \in \mathbb{Z}^n_+$ be an integer lattice vector with positive entries. It defines the *weight function* $\omega^\star : Q \to \mathbb{R}$ on $Q$ by $\omega^\star(v) = \langle \omega, v \rangle$, where $\langle , \rangle$ is the standard scalar product in $\mathbb{R}^n$. Note that $Q$ lies in a hyperplane whose normal vector is $(1, \ldots, 1)$. The weight function $\omega^\star$ is maximized along a face $G_\omega$ of $Q$ which is uniquely determined by the condition that the vector $\omega$ lies in the relative interior of its normal cone $\omega \in \sigma^\circ_{G_\omega}$. A weight function $\omega^\star$ is called $Q$-*generic* if $G_\omega$ is a vertex of $Q$.

If $Q = Pe^{n-1}$ is the standard permutohedron the previous consideration shows that each $\omega \in \mathbb{Z}^n_+$ determines a unique face $\mathcal{F}_\omega$ by the condition $\mathcal{F} = \mathcal{F}_\omega$ if and only if $\omega \in \sigma^\circ_{\mathcal{F}}$.

**Definition 4.2.** For a generalized permutohedron $Q$ let

$$F_q(Q) = \sum_{\omega \in \mathbb{Z}^n_+} q^{\mathrm{rk}_Q(\mathcal{F}_\omega)} \mathbf{x}_\omega.$$

For a face $\mathcal{F}$ of $Pe^{n-1}$ the enumerator of integer vectors $\omega \in \mathbb{Z}^n_+$ such that $\mathcal{F} = \mathcal{F}_\omega$ is exactly the monomial quasisymmetric function

$$M_{\mathrm{type}(\mathcal{F})} = \sum_{\omega \in \sigma^\circ_{\mathcal{F}}} \mathbf{x}_\omega.$$

We obtain the expansion of $F_q(Q)$ in the monomial basis

$$F_q(Q) = \sum_{\mathcal{F} \in L(Pe^{n-1})} q^{\mathrm{rk}_Q(\mathcal{F})} M_{\mathrm{type}(\mathcal{F})}. \tag{3}$$

By definition of the map $\pi_Q$ we have

$$\sum_{\omega\in\sigma_G^\circ}\mathbf{x}_\omega=\sum_{\mathcal{F}:\pi_Q(\mathcal{F})=G}M_{\mathrm{type}(\mathcal{F})},$$

which gives the expansion of $F_q(Q)$ in terms of the face lattice of $Q$

$$F_q(Q)=\sum_{G\in L(Q)}q^{\dim(G)}\sum_{\mathcal{F}:\pi_Q(\mathcal{F})=G}M_{\mathrm{type}(\mathcal{F})}. \tag{4}$$

From this representation we easily derive that the enumerator $F_q$ contains the information about $f$-vectors of generalized permutohedra. Recall that the $f$-vector of a convex $(n-1)$-dimensional polytope $P$ is the integer vector $f=(f_0,f_1,\ldots,f_{n-1})$, where $f_i$ is the number of $i$-dimensional faces of $P$. It is codified by the $f$-polynomial

$$f(P,q)=f_0+f_1q+f_2q^2+\cdots+f_{n-1}q^{n-1}.$$

The following theorem is a common generalization of [10, Theorem 1.1] and [12, Theorem 1] which are proven by using particular combinatorial properties of graphical zonotopes and nestohedra.

**Theorem 4.3.** *The $f$-polynomial $f(Q,q)$ of an $(n-1)$-dimensional generalized permutohedron $Q$ is determined by the principal specialization*

$$f(Q,q)=(-1)^n\mathbf{ps}(F_{-q}(Q))(-1).$$

*Proof.* By formula (4) and the fact $\mathbf{ps}(M_{\mathrm{type}(\mathcal{F})})(-1)=(-1)^{|\mathcal{F}|+1}$ the principal specialization of $F_q(Q)$ gives

$$(-1)^n\mathbf{ps}(F_{-q}(Q))(-1)=\sum_{G\in L(Q)}q^{\dim(G)}\sum_{\mathcal{F}:\pi_Q(\mathcal{F})=G}(-1)^{|\mathcal{F}|+1+n+\dim(G)}.$$

The inner sum is equal to 1 by Proposition 2.2, which completes the proof. □

We review some specializations of the quasisymmetric function $F_q(Q)$. By taking $q=0$ it is specified to the Billera, Jia and Reiner quasisymmetric function $F(Q)$ which is the enumerator of $Q$-generic weight vectors

$$F(Q)=\sum_{\omega\in\mathbb{Z}_+^n:\mathrm{rk}(\mathcal{F}_\omega)=0}\mathbf{x}_\omega.$$

Theorem 4.3 may be regarded as the generalization of the formula for number of vertices $f_0$ of a generalized permutohedron $Q$, see [3, Theorem 9.2]

$$f_0 = (-1)^n \mathbf{ps}(F(Q))(-1).$$

Note that $F_q(Q)$ degenerates at $q = 1$ since $F_1(Q) = \sum_{\omega \in \mathbb{Z}^n_+} \mathbf{x}_\omega = (M_1)^n$.

For particular classes of generalized permutohedra the enumerator $F_q$ specializes to known quasisymmetric invariants as it is announced in introduction. We briefly review these cases.

**Grafical zonotopes** The vertices of a graphical zonotope $Z_\Gamma$ are in one-to-one correspondence with regions of the corresponding graphical arrangement. The integer points $\omega \in \mathbb{Z}^n_+$ can be interpreted as graph colorings by labelling vertices of a graph $\Gamma$. The normal cone at a vertex of $Z_\Gamma$ is a region of the corresponding graphical arrangements. So $\omega$ determines a proper coloring of the graph $\Gamma$ if and only if $\mathrm{rk}_{Z_\Gamma}(\mathcal{F}_\omega) = 0$. Thus the enumerators of proper colorings and $Z_\Gamma$-generic integer vectors are equal, i.e. $F_0(Z_\Gamma) = X_\Gamma$, where $X_\Gamma$ is the Stanley chromatic symmetric function (introduced in [18]).

The main argument in [11] in the proof of Theorem 4.3 for graphical zonotopes was based on the Humpert and Martin cancelation-free formula for the antipode of the chromatic Hopf algebra of graphs [13].

**Nestohedra** The proof of Theorem 4.3 in the case of nestohedra given in [12] is based on the recurrence relation satisfied by $f$-polynomials of nestohedra obtained by Postnikov in [15].

# 5 Matroid base polytope

A standard reference monograph for matroid theory is [14]. The most adequate for our purposes is a definition of a matroid as a collection of bases. A matroid $M = ([n], \mathcal{B})$ on the ground set $[n]$ is a nonempty collection $\mathcal{B}$ of subsets of $[n]$ such that the exchange property is satisfied:

- For each $B_1, B_2 \in \mathcal{B}$ and $i \in B_1 \setminus B_2$ there is $j \in B_2 \setminus B_1$ such that $B_1 \setminus \{i\} \cup \{j\} \in \mathcal{B}$.

The members of $\mathcal{B}$ are called bases of $M$ and we denote the collection of bases of $M$ by $\mathcal{B}(M)$. All bases have the same number of elements which is called the rank $r(M)$ of the matroid $M$.

**Definition 5.1.** The matroid base polytope is a convex polytope

$$P_M = \mathrm{Conv}\left\{e_B \mid B \in \mathcal{B}(M)\right\},$$

where $e_B = \sum_{i \in B} e_i$ and $e_i, i = 1, \ldots, n$ are the standard basis vectors in $\mathbb{R}^n$.

The polytope $P_M$ is contained in the hypersimplex

$$\Delta_r^n = [0,1]^n \cap \{x_1 + x_2 + \cdots + x_n = r(M)\}.$$

The following characterization of matroids in terms of their base polytopes, proven in [7, Section 2.2, Theorem 1] and [8, Theorem 4.1] shows that matroid base polytopes are generalized permutohedra.

**Theorem 5.2.** *Let $\mathcal{S}$ be a collection of $r$-subsets of $[n]$ and $P_{\mathcal{S}}$ be a convex polytope in $\mathbb{R}^n$*

$$P_{\mathcal{S}} = \mathrm{Conv}\left\{e_S \mid S \in \mathcal{S}\right\},$$

*where $e_S = \sum_{i \in S} e_i$. Then $\mathcal{S}$ is the collection of bases of a matroid if and only if every edge of $P_{\mathcal{S}}$ is a translate of the vector $e_i - e_j$, for some $i, j \in [n]$.*

**Definition 5.3.** Given a matroid $M = ([n], \mathcal{B})$ the elements $i, j \in [n]$ are equivalent if there exist bases $B_1$ and $B_2$ with $i \in B_1$ and $B_2 = (B_1 \backslash \{i\}) \cup \{j\}$. The equivalence classes are called *connected components* of $M$. The matroid $M$ is *connected* if it has only one connected component.

Denote by $c(M)$ the number of connected components of a matroid $M$.

**Proposition 5.4** ([6], Proposition 2.4)**.** *The dimension of the matroid base polytope $P_M$ corresponding to a matroid $M$ on $[n]$ is determined by*

$$\dim(P_M) = n - c(M).$$

The dual matroid is defined by the set of bases

$$\mathcal{B}(M^*) = \{[n] \setminus B \mid B \in \mathcal{B}(M)\}.$$

The affine transformation of the $n$-cube $x \mapsto \mathbf{1} - x, x \in [0,1]^n$, where $\mathbf{1} = (1, \ldots, 1)$ determines an affine isomorphism between $P_M$ and $P_{M^*}$.

## 5.1 Flags of matroid base polytope

For a vector $\omega = (\omega_1, \omega_2, \ldots, \omega_n) \in \mathbb{Z}_+^n$ we regard the function $\omega^*$ on the matroid base polytope $P_M$ as a weight function on a matroid $M$, so that the $\omega$-weight of a basis $B = \{i_1, \ldots, i_{r(M)}\} \in \mathcal{B}(M)$ is given by

$$\omega^*(B) = \langle \omega, e_B \rangle = \omega_{i_1} + \cdots + \omega_{i_{r(M)}}.$$

Let $\mathcal{B}_\omega \subset \mathcal{B}(M)$ be the collection of bases of $M$ having maximum $\omega$-weight. The collection $\mathcal{B}_\omega$ consists of all bases obtained by greedy algorithm. The polytope $P_{\mathcal{B}_\omega}$ is the face of $P_M$ on which $\omega^*$ is maximized. Theorem 5.2 implies that $\mathcal{B}_\omega$ is the collection of bases of a matroid which we denote by $M_\omega$.

Recall that $\omega$ determines a unique flag of subsets

$$\mathcal{F}_\omega : \emptyset =: F_0 \subset F_1 \subset \ldots \subset F_k \subset F_{k+1} := [n] \tag{5}$$

for which $\omega$ is constant on $F_i \setminus F_{i-1}, i = 1, \ldots, k+1$ and satisfies $\omega|_{F_{i+1} \setminus F_i} < \omega|_{F_i \setminus F_{i-1}}, i = 1, \ldots k$. The following proposition shows that $M_\omega$ depends only on the flag $\mathcal{F}_\omega$. By $r(A) = r(M\mid_A)$, where $M\mid_A$ is the restriction of $M$ to $A \subset [n]$ is defined the rank function of $M$.

**Proposition 5.5** ([2], Proposition 1)**.** *Let $\omega \in \mathbb{Z}_+^n$ and $\mathcal{F} = \mathcal{F}_\omega$ is the corresponding flag given by* (5)*. A base $B \in \mathcal{B}(M)$ of a matroid $M$ has the maximum $\omega$-weight, i.e. $B \in B_\omega$ if and only if $|B \cap (F_i \setminus F_{i-1})| = r(F_i) - r(F_{i-1}), i = 1, \ldots, k+1$.*

We denote the matroid corresponding to a flag $\mathcal{F}$ by $M_\mathcal{F}$. It determines the map $\pi_{P_M} : L(Pe^{n-1}) \to L(P_M)$ associated to the matroid base polytope, namely $\pi_{P_M}(\mathcal{F}) = P_{M_\mathcal{F}}$. The following proposition describes the matroid $M_\mathcal{F}$.

**Proposition 5.6** ([2], Proposition 2)**.** *If $\mathcal{F} = \{\emptyset =: F_0 \subset \ldots \subset F_{k+1} := [n]\}$, then*

$$M_\mathcal{F} = \bigoplus_{i=1}^{k+1} \left(M|_{F_i}\right) / F_{i-1}.$$

As it holds the following property for the direct sum of matroids $P_{M_1 \oplus M_2} = P_{M_1} \times P_{M_2}$, we obtain

**Corollary 5.7.** *The $P_M$-rank of a flag $\mathcal{F}$ is given by*

$$\mathrm{rk}_{P_M}(\mathcal{F}) = \dim(P_{M_\mathcal{F}}) = n - \sum_{i=1}^{k} c\left((M|F_i)/F_{i-1}\right).$$

## 5.2 Quasisymmetric enumerator $F_q(P_M)$ for matroids

The Hopf algebra of matroids $\mathcal{M}at$ was introduced by Schmitt [17] and more intensively studied by Crapo and Schmitt [4], [5].

As a vector space $\mathcal{M}at$ is linearly spanned over the field $\mathbf{k}$ by isomorphism classes of finite matroids $[M]$. The direct sum induces a product, while a coproduct is determined by restrictions and contractions

$$[M_1] \cdot [M_2] = [M_1 \oplus M_2] \quad \text{and} \quad \Delta[M] = \sum_{A \subseteq E} [M|_A] \otimes [M/A].$$

The Hopf algebra $\mathcal{M}at$ is graded $\mathcal{M}at = \bigoplus_{n \geq 0} \mathcal{M}at_n$, where $\mathcal{M}at_n$ denotes the subspace spanned by elements $[M]$ for which the ground set has cardinality $n$. The Hopf algebra $\mathcal{M}at$ is graded, connected commutative, but not-cocommutative. The unit is the class of a unique matroid on the empty set $[M_\emptyset]$.

The character $\zeta : \mathcal{M}at \to \mathbf{k}$ defined by

$$\zeta([M]) = \begin{cases} 1, & M \text{ is a direct sum of loops and isthmuses,} \\ 0, & \text{otherwise} \end{cases}$$

turns $\mathcal{M}at$ into a combinatorial Hopf algebra considered by Billera, Jia and Reiner in [3]. By Theorem 3.1 there is a unique morphism $\Psi : (\mathcal{M}at, \zeta) \to (QSym, \zeta_Q)$ of combinatorial Hopf algebras given in monomial bases of quasisymmetric functions by

$$\Psi([M]) = \sum_{\alpha \models n} c_\alpha([M]) M_\alpha.$$

**Proposition 5.8** ([3], Proposition 3.3)**.** *The coefficient $c_\alpha([M])$ is the number of flags $\mathcal{F} : \emptyset = F_0 \subset \cdots \subset F_{k+1} = [n]$ having* $\mathrm{type}(\mathcal{F}) = \alpha$ *and for which each subquotient $(M|_{F_i})/F_{i-1}$ is a direct sum of loops and isthmuses.*

We extend the basic field $\mathbf{k}$ into the field of rational functions $\mathbf{k}(q)$ and define the character $\zeta_q : \mathcal{M}at \to \mathbf{k}(q)$ with

$$\zeta_q([M]) = q^{\mathrm{rk}(M)},$$

where $\mathrm{rk}(M) = n - c(M)$. Let $\Psi_q : (\mathcal{M}at, \zeta_q) \to (\mathcal{Q}Sym, \zeta_\mathcal{Q})$ be a unique morphism of combinatorial Hopf algebras over $\mathbf{k}(q)$.

**Proposition 5.9.** *The quasisymmetric enumerator $F_q(P_M)$ for a matroid base polytope is determined by*

$$F_q(P_M) = \Psi_q([M]).$$

*Proof.* The morphism $\Psi_q$ is given by

$$\Psi_q([M]) = \sum_{\alpha \models n} c_{q,\alpha}([M]) M_\alpha,$$

where

$$c_{q,\alpha}([M]) = \sum_{\mathcal{F}:\mathrm{type}(\mathcal{F})=\alpha} \prod_{i=1}^{k} q^{\mathrm{rk}\left((M|_{F_i})/F_{i-1}\right)}.$$

By Corollary 5.7 we have

$$c_{q,\alpha}([M]) = \sum_{\mathcal{F}:\mathrm{type}(\mathcal{F})=\alpha} q^{\mathrm{rk}_{P_M}(\mathcal{F})}.$$

Consequently,

$$\Psi_q([M]) = \sum_{\mathcal{F}} q^{\mathrm{rk}_{P_M}(\mathcal{F})} M_{\mathrm{type}(\mathcal{F})},$$

where the sum is over all flags of $[n]$. This is exactly the form of $F_q(P_M)$ given by identity (3). □

As a consequence of the previous proposition and theorem 4.3 we obtain the following algebraic description of the $f$-polynomials of matroid base polytopes.

**Theorem 5.10.** *The $f$-polynomial of a matroid base polytope $P_M$ is given by*

$$f(P_M, q) = (-1)^n \mathbf{ps}(\Psi_{-q}([M]))(-1).$$

## 5.3 The uniform matroid base polytope

We use theorem 5.10 to calculate $f$-polynomials of uniform matroid base polytopes $P_{U_{n,r}}$. The uniform matroid $U_{n,r}$ is a matroid with the set of bases

$\mathcal{B}(U_{n,r}) = \binom{[n]}{r}$ consisting of all $r$-elements subsets of $[n]$. It follows from definition that $P_{U_{n,r}}$ is the hypersimplex

$$P_{U_{n,r}} = \Delta^n_r.$$

Note that $\mathcal{B}(U_{n,0}) = \{\emptyset\}$ and $\mathcal{B}(U_{n,n}) = \{[n]\}$, so $P_{U_{n,0}} = \mathbf{0}$ and $P_{U_{n,n}} = \mathbf{1}$ are single points, where $\mathbf{0} = (0, \ldots, 0)$ and $\mathbf{1} = (1, \ldots, 1)$.

Consider the uniform matroid $U_{n,r}$ and assume that $0 < r < n$. Let $\mathcal{F} : \emptyset = F_0 \subset F_1 \subset \ldots \subset F_{k+1} = [n]$ be a flag with $\text{type}(\mathcal{F}) = (\alpha_1, \ldots, \alpha_i, \alpha_{i+1}, \ldots, \alpha_k)$. Define $i_0 = i_0(\mathcal{F}) = \min\{i \mid \alpha_1 + \cdots + \alpha_i \geq r\}$ and partial sums $r' = \alpha_1 + \cdots + \alpha_{i_0-1}, r'' = r' + \alpha_{i_0}$. By proposition 5.6 we find

$$M_{\mathcal{F}} = U_{r',r'} \oplus U_{\alpha_{i_0}, r-r'} \oplus U_{n-r'',0}.$$

It shows that faces of the uniform matroid base polytopes are uniform matroid base polytopes as well, the fact that is obvious from the description by hypersimplices. We have $P_{M_{\mathcal{F}}} \cong P_{U_{\alpha_{i_0}, r-r'}}$, which implies the following formula for the $P_M$-rank, where $M = U_{n,r}$

$$\text{rk}_{P_M}(\mathcal{F}) = \dim(P_{M_{\mathcal{F}}}) = \begin{cases} 0, & r = r'', \\ \alpha_{i_0} - 1, & r < r''. \end{cases} \tag{6}$$

Denote the concatenation of monomial quasisymmetric functions by

$$M_\alpha \circ M_\beta = M_{\alpha\beta}.$$

We have

$$F_q(P_{U_{n,r}}) = \sum_{\mathcal{F}:\text{rk}_{P_M}(\mathcal{F})=0} M_{\text{type}(\mathcal{F})} + \sum_{\mathcal{F}:\text{rk}_{P_M}(\mathcal{F})>0} q^{\alpha_{i_0(\mathcal{F})-1}} M_{\text{type}(\mathcal{F})}.$$

Finally,

$$F_q(P_{U_{n,r}}) = \binom{n}{r} (M_1)^r \circ (M_1)^{n-r} +$$

$$+ \sum_{0 \leq r' < r < r' + \lambda \leq n} \binom{n}{r'} \binom{n-r'}{\lambda} q^{\lambda - 1} (M_1)^{r'} \circ M_\lambda \circ (M_1)^{n-r'-\lambda}.$$

The principal specilization evaluated at $-1$ gives

$$f(P_{U_{n,r}},q)=\binom{n}{r}+\sum_{0\leq r'<r<r'+\lambda\leq n}\binom{n}{r'}\binom{n-r'}{\lambda}q^{\lambda-1}.$$

It determines the $f$-vector of $P_{U_{n,r}}$ by $f_0=\binom{n}{r}$ and in terms of trinomial coefficients

$$f_{k-1}=\sum_{0\leq r'<r<r'+k\leq n}\binom{n}{r',k,n-r'-k},\ \text{ for }\ k=2,\ldots,n.$$

**Example 5.11.** The polytope $P_{U_{4,2}}=\Delta_2^4$ is an octahedron. We obtain

$$F_q(P_{U_{4,2}})=6\left(M_{(2,2)}+2M_{(1,1,2)}+2M_{(2,1,1)}\right)q^0+12M_{(1,2,1)}q+$$

$$+4\left(M_{(1,3)}+M_{(3,1)}\right)q^2+M_{(4)}q^3,$$

which yields to the expected expression $f(P_{U_{4,2}},q)=6+12q+8q^2+q^3$.

# References


[1] M. Aguiar, N. Bergeron, F. Sottile, Combinatorial Hopf algebras and generalized Dehn-Sommerville relations, Compositio Mathematica 142 (2006) 1–30

[2] F. Ardila, C. Klivans, The Bergman complex of a matroid and phylogenetic trees, J. Combin. Theory Ser. B 96 (2006) 38–49.

[3] L. Billera, N. Jia, V. Reiner, A quassisymetric function for matroids, European J. Comb. 30 (2009) 1727–1757

[4] H. Crapo, W. Schmitt, A free subalgebra of the algebra of matroids, European J. Combin. 26 (2005) 1066–1085.

[5] H. Crapo, W. Schmitt, A unique factorization theorem for matroids, J. Combin. Theory Ser. A 112 (2005) 222–249.

[6] E. M. Feichtner, B. Sturmfels, Matroid polytopes, nested sets and Bergman fans, Port. Math. (N.S.) 62 (2005) no.4, 437-468

[7] I. M. Gel'fand, V. V. Serganova, Combinatorial geometries and torus strata on homogeneous compact manifolds, Russian Math. Surveys 42 (2) (1987) 133–168.

[8] I. M. Gel'fand, M. Goresky, R. MacPherson, V. Serganova, Combinatorial geometries, convex polyhedra, and Schubert cells, Adv. Math. 63 (1987) 301-316.

[9] D. Grinberg, V. Reiner, Hopf Algebras in Combinatorcs, arXiv:1409.8356

[10] V. Grujić, Quasisymmetric functions for nestohedra, arXiv:1409.1420

[11] V. Grujić, Counting faces of graphical zonotopes, Ars Math. Contemporanea, 13 (2017) 227–234.

[12] V. Grujić, T. Stojadinović, Counting faces of nestohedra, arXiv:1703.08826

[13] B. Humpert and J. Martin, The incidence Hopf algebra of graphs, *SIAM J. Discrete Math.*, **26** (2) (2012), 555–570.

[14] J. G. Oxley, Matroid Theory, Oxford University Press, New York, 1992.

[15] A. Postnikov, Permutohedra, associahedra, and beyond, Int. Math. Res. Notices 2009 (2009) 1026–1106

[16] A. Postnikov, V. Reiner, L. Williams, Faces of generalized permutohedra, Documenta Math. 13 (2008) 207–273

[17] W. R. Schmitt. Incidence Hopf algebras, J. Pure Appl. Algebra 96 (1994), no. 3, 299-330.

[18] R. P. Stanley, A symmetric function generalization of the chromatic polynomial of a graph, Adv. Math. 111 (1995), 166–194.